

IAI SPECIAL EDITION

RESEARCH ARTICLE

Antibiotic use and antibiotic resistance profile of bacteria isolated from out-patients of Pakem primary health care Yogyakarta

Daru Estiningsih^{1,2}, Ika Puspitasari^{1,3}, Titik Nuryastuti⁴, Endang Lukitaningsih¹

¹ Faculty of Pharmacy, Universitas Gadjah Mada, Yogyakarta, Indonesia

² Department of Pharmacy, Faculty of Health Sciences, Universitas Alma Ata, Yogyakarta, Indonesia

³ Academic Hospital, Universitas Gadjah Mada, Yogyakarta, Indonesia

⁴ Faculty of Medicine, Public Health, and Nursing Universitas Gadjah Mada, Yogyakarta, Indonesia

Keywords

Antibiotic resistance
Bacteria
ESCAPE organisms
Infection

Correspondence

Ika Puspitasari
Faculty of Pharmacy
Universitas Gadjah Mada
Yogyakarta
Indonesia
ika.puspitasari@gmail.com

Abstract

Background: The use of antibiotics is one of the procedures in the treatment of infectious diseases. Antibiotic are given to kill or inhibit the growth of infectious bacteria within the host's system. Inappropriate use of antibiotics may lead to antibiotic resistance where the bacteria able to resist the effect of the drugs. **Objective:** The purpose of this study was to determine the pattern of antibiotic use and antibiotic resistance in out-patients at the Pakem Health Center, Yogyakarta. **Method:** This study employed a descriptive research design using the data of patients with bacterial infections. Twenty-six patients were the subjects of the study with predetermined characteristics, namely age, gender, occupation, and antibiotics given. **Result:** The result showed the five (5) common types of infections were pyoderma, pharyngitis, abscess, bronchitis, and wound infection. Meanwhile, the infectious bacteria found were *Staphylococcus aureus*, *Streptococcus pyogenes*, *Klebsiela pneumoniae*, *Acinetobacter baumannii*, *Klebsiella oxytoca*, *Pseudomonas aeruginosa*, and *Streptococcus betahaemolyticus*. The prescribed antibiotics included amoxicilin (73.33%), metronidazole (16.67%), gentamicin (6.67%), and ciprofloxacin (3.33%). The antibiotic resistant bacteria were *Acinetobacter baumannii*, *Klebsiela pneumoniae*, *Klebsiella oxytoca*, and *Staphylococcus aureus*. Among the test bacteria, several groups showed multiple-drug resistance. The five highest antibiotic resistance found were penicillin (52.9%), ampicillin (40.0%), tetracycline (33.3%), cefuroxime (33.3%), and amikacin (33.3%). **Conclusion:** The most common type of infection was pyoderma and the antibiotic treatment given to most occurring cases of infection was amoxicillin.

Introduction

One of the procedures to handle infectious diseases is antimicrobial therapy, which aims to kill or inhibit the growth of infectious bacteria in the body (Kemenkes, 2017; Ho & Ip, 2019). More than 90% of cases of respiratory tract infection and pharyngitis were administered antibiotics which resulted in an increased number of antibiotic prescriptions (Apsari *et al.*, 2017). Moreover, in America, the use of antibiotics increased among adult patients (about 192-198 million) in 2014 (Ho & Ip, 2019).

Antibiotics are powerful medicines that are taken to kill or inhibit the growth of bacteria in the body. However, the excessive use of antibiotics may trigger the emergence of antibiotic resistance (Humaida, 2014). Excessive use of antibiotics has an impact not only in terms of resistance but also affects the level of effectiveness and efficiency (Friedman *et al.*, 2016). This is because it is associated with increased morbidity, mortality, length of treatment, and care costs (Sinto, 2020). The resistant bacteria tend to cause the failure of infectious disease therapy because they have become resistant to antibiotics (Mahdani *et al.*,

2020). Antibiotic resistance in ESCAPE organisms (*Enterococcus*, *Staphylococcus aureus*, *Klebsiella pneumoniae*, *Acinetobacter baumannii*, *Pseudomonas aeruginosa*, and *Enterobacter* sp) is frequently linked to significantly higher rates of morbidity, mortality, and fiscal burden (Zhen et al., 2019).

Antibiotic resistance has become a major global problem (Murray et al., 2022). In 2009, Indonesia was ranked 8th out of 27 countries with the highest multidrug-resistant predicate in the world (Estiningsih et al., 2016). Bacteria can be resistant to an antibiotic due to several factors, such as the lack of cell organelles that are the target of antibiotics, certain bacterial species having cell walls which are not permeable to certain antibiotics, and a population of bacterial species do not necessarily have the same level of sensitivity to an antibiotic (Kemenkes, 2017). One way to lower the incidence of antibiotic resistance is to reduce the number of antibiotic prescriptions (Apsari et al., 2017).

Appropriate use of antibiotics may reduce adverse events. The differences in the resistance patterns of bacteria determine the antibiotic prescription. Hence, this study aims to determine the antibiotic use and antibiotic resistance patterns of infectious bacteria associated with the out-patients within the primary health care at Pakem, to aid the effectiveness and efficiency of antibiotic prescription.

Methods

Design

This research employed a descriptive study with prospective data collection carried out by analysing patients' medical record data obtained from the Pakem primary health care, Yogyakarta.

Materials

The instrument used in this study was the research data

form. The material used was the medical records of the patients and the specimen taken from patients according to the infection.

Inclusion and exclusion criteria

The inclusion criteria included out-patients in Pakem primary health care, Yogyakarta, who were diagnosed with infections and received antibiotic therapies, patients who were not undergoing haemodialysis, patients who were not treated from the hospital for at least 30 days, and patients that were willing to become probandi in this study. Meanwhile, the exclusion criterion was vegetarian patients.

Results

A total of 26 patient data were analysed. Patients who participated in the study were dominated by those in the late adolescent age (30.8%), followed by the early adolescents (15.4%), late adults (15.4%), early elderly (15.4%), late elderly (15.4%) and the early adults (7.7%) as shown in Table I.

Table I: Age characteristics of patients

Age	n	%
Early adolescence (12 - 16 years old)	4	15.4
Late adolescence (17 - 25 years old)	8	30.8
Early adult (26 - 35 years old)	2	7.7
Late adult (36- 45 years old)	4	15.4
Early elderly (46 - 55 years old)	4	15.4
Late elderly (56 - 65 years old)	4	15.4
Total	26	100

The results of the study obtained from the 26 patients showed various types of infections, most of which were pyoderma (19.3%), pharyngitis (15.4%), abscess (11.5%), bronchitis (11.5%), wound infectious (7.7%), otitis media (7.69%), cellulitis (3.9%), and sinusitis (3.9%) (Figure 1).




Figure 1: Percentage of infection type

Infections were caused by several types of bacteria with the largest percentage in *Staphylococcus aureus* (45.8%) followed by *Streptococcus pyogenes* (25%), *Klebsiella pneumonia* (12.5%), *A.baumannii* (4.2%), *Klebsiella oxytoca* (4.2%), *P.aeruginosa* (4.2%) and *Streptococcus betahaemolyticus* (4.2%) (Table II).

Table II: Bacteria causing infections

Bacteria	Total	%
<i>Staphylococcus aureus</i>	11	45.8
<i>Streptococcus pyogenes</i>	6	25.0
<i>Klebsiella pneumoniae</i>	3	12.5
<i>Pseudomonas aeruginosa</i>	1	4.2
<i>Acinetobacter baumannii</i>	1	4.2
<i>Streptococcus betahaemolyticus</i>	1	4.2
<i>Klebsiella oxytoca</i>	1	4.2
Total	24	100

In this study, data on the use of antibiotics for treatment were also obtained. The data showed that the most widely utilised antibiotics were amoxicillin 500 mg tablets (73.3%), metronidazole tablets (16.7%), then gentamicin (6.7%), and ciprofloxacin 500 mg tablets (3.3%) (Table III). Most amoxicillin was prescribed to infectious patients, especially those diagnosed with pharyngitis, followed by other types of infections such as

dermatitis, abscesses, infectious wounds, and otitis media. Amoxicillin is a β -lactam antibiotic that has a broad spectrum and it is often used to treat various infectious diseases caused even by gram-positive and gram-negative bacteria, such as ear infections, pneumonia, streptococcal pharyngitis, skin infections, urinary tract infections, Salmonella infections, Chlamydia infections and Lyme disease (Benjamin, 2019). Amoxicillin is also a type of antibiotic that has been listed in the Indonesian National Formulary so that patients are guaranteed to get the right, efficacious, quality, safe, and affordable medicines (Apsari et al., 2017).

Table III: Antibiotic use

Antibiotic	Total	%
Amoxicillin	22	73.3
Metronidazole	5	16.7
Gentamicin	2	6.7
Ciprofloxacin	1	3.3
Total	30	100

Based on the patient’s data, antimicrobial susceptibility testing (AST) was then carried out to determine antibiotic sensitivity. The result revealed the diversity in the bacterial resistance pattern to the test antibiotics (Figure 2; Table IV).




Figure 2: Antibiotic resistance (%)

Table IV: Antibiotic sensitivity test results

Group	Antibiotic	Bacteria (% resistant)						
		A	B	C	D	E	F	G
Aminoglycosides	Gentamicin	100	0	0	0	33.3	-	-
	Tobramycin	0	0	0	0	-	-	-
	Amikacin	100	33.3	0	0	-	-	-
B-lactam + β lactamase inhibitor	Ampi-sulbactam	0	-	-	-	-	-	-
	Amoxiclav	-	0	0	-	-	-	-
Piperacillin + tazobactam	Tazobactapiperacillin	0	0	0	0	-	-	-
Cephalosporin	Cefazolin	-	0	100	-	0	-	-
	Cefuroxime	-	33.3	0	-	-	-	-
	Cefepime	0	0	0	0	-	0	0
	Ceftriaxone	100	0	0	-	-	0	0
	Ceftazidime	0	-	-	0	-	-	-
	Cefotaxime	100	-	-	-	-	0	0
Fluoroquinolone	Ciprofloxacin	0	0	0	0	9.1	-	-
	Levofloxacin	0	0	0	0	-	0	0
Carbapenem	Meropenem	0	0	0	0	-	-	-
Cotrimoxazole	Cotrimoxazole	0	0	0	-	0	-	-
Monobactam	Aztreonam	-	-	-	0	-	-	-
Penicillin	Ampicillin	-	100	100	-	-	0	0
	Piperacillin	-	-	-	0	-	-	-
	Penicillin	-	-	-	-	81.8	0	0
Macrolides	Erythromycin	-	-	-	-	0	0	0
Oxazolidinone	Linezolid	-	-	-	-	0	-	-
Tetracycline	Tetracycline	-	-	-	-	36.4	-	0
Lincosamide	Clindamycin	-	-	-	-	0	0	0
Glycopeptide	Vancomycin	-	-	-	-	-	0	0
Chloramphenicol	Chloramphenicol	-	-	-	-	25.0	0	0

Note: A: (*Acinetobacter baumannii*); B: (*Klebsiella pneumoniae*); C: (*Klebsiella oxytoca*); D: (*Pseudomonas aeruginosa*); E: (*Staphylococcus aureus*); F: (*Streptococcus betahaemolyticus*); G: (*Streptococcus pyogenes*)

Discussion

Based on the result of the antibiotic sensitivity test, *A. baumannii* showed 100% resistance to gentamicin and amikacin (aminoglycosides). The most studied mechanism of resistance to aminoglycoside antibiotics in strain *A. baumannii* is the production of aminoglycositic-modifying enzymes (AMEs) (Kyriakidis et al., 2021). In this study, *A. baumannii* also showed absolute resistance to ceftriaxone (100%) and Cefotaxime (100%). Additionally, based on the result of antibiotic susceptibility testing (AST) between 1995 and 2004, it was also confirmed that there was an increase in the percentage of resistance to all antibiotics in four main classes, namely fluoroquinolones (50-73%), aminoglycosides (19-31%), β -lactam (39-66%), and carbapenems (9-39%) (Vázquez-López et al., 2020). This indicates that the bacteria have developed resistance to survive.

Klebsiella pneumoniae showed similar low resistance to amikacin and cefuroxime (33.3%), while to Penicillin by 100%. Moreover, it is also reported in other studies where most of the *K. pneumoniae* isolates were

classified as multidrug-resistant (MDR) with high levels of resistance to a broad spectrum of drugs including β -lactams, aminoglycosides, and quinolones (Fair & Tor, 2014). Antibiotics from the β -lactam group are commonly prescribed worldwide including penicillins, cephalosporins, monobactams, and carbapenems (Ur Rahman et al., 2018). The resistance mechanism of β -lactam antibiotics-resistant bacteria is the production of β -lactamase enzymes in the presence of cell wall transpeptides (Igrejas et al., 2019).

Klebsiella oxytoca showed 100% resistance to cefazolin and ampicillin. Other studies also mentioned the presence of *K. oxytoca* carrying a constitutive β -lactamase low resistance to penicillin but susceptible to other β -lactams (Fenosa et al., 2009). Furthermore, *Staphylococcus aureus* was resistant to several antibiotics and the largest percentage was to a penicillin (81.8%). *S. aureus* resistance to penicillin is associated with Methicillin-Resistant *Staphylococcus Aureus* (MRSA) (Laux et al., 2019). It can produce penicillinase which can rehydrate the β -lactam ring of penicillin, causing the antibiotic to be insensitive. MRSA is a kind of multi-drug resistant "super bacteria" that

are resistant to penicillin, cephalosporin, chloramphenicol, lincomycin, aminoglycosides, tetracycline, macrolides, quinolones, sulfonamides and rifampicin. Such is a very difficult problem in clinical medicine (Guo *et al.*, 2020). In addition, *S. aureus* showed resistance to tetracycline (36.4%), gentamicin (33.3%), chloramphenicol (25%) and ciprofloxacin (9.1%). Meanwhile, *P. aeruginosa*, *Streptococcus betahaemolyticus*, and *Streptococcus pyogenes* were not resistant to any antibiotics in this study.

The result of the sensitivity test showed that several bacteria in this study belong to the multidrug-resistant organisms (MDRO) category (Table III). MDRO is a microorganism, especially bacteria that show resistance to at least one type of antibiotic from in ≥ 3 antibiotic group (Siegel *et al.*, 2017). This study showed that *Staphylococcus aureus* showed resistance to five (n=5) groups of antibiotics, one of which was the Penicillin group (Penicillin). This resistance might be due to the production of penicillinase which is in charge of degrading penicillin in the extracellular space by hydrolyzing the ring β -lactam (Ho & Ip, 2019). In addition, *Klebsiella pneumoniae* was also resistant to three (n=3) types of antibiotic groups, namely aminoglycosides (amikacin), cephalosporins (cefuroxime), and penicillin (penicillin).

In most cases, infections caused by resistant bacteria result in twice higher adverse outcomes as those caused by susceptible bacteria (Cosgrove & Carmeli, 2003). These negative effects could be clinical (death or therapy failure) or economic (cost of treatment or length of stay). Additionally, it makes a note of delayed therapy and antibiotic treatment failures to eradicate the infection (Friedman *et al.*, 2016). The published MDROs restrain guidance remarks ongoing debate areas on optimal control strategies (Siegel *et al.*, 2017). Moreover, various approaches have diminished MDROs rates. Thus, intervention options for controlling MDRO transmission must be based on local problem assessments, the prevalence of various MDROs and feasibility (Siegel *et al.*, 2017). Furthermore, individuals ought to seek proper guidance as well as adopt effective measures that care for their needs and circumstances (Siegel *et al.*, 2017). The management of patients infected with resistant bacteria is conducted by providing appropriate antibiotics; considering their clinical efficacy, sensitivity, patient clinical condition, cost, prioritizing first-line/narrow-spectrum antibiotics, and their availability in healthcare facilities (Kemenkes, 2011)

This study showed that most of the isolates resistant to antibiotics were gram-negative where *A. baumannii* had the highest percentage (30.8%), followed by *Staphylococcus aureus* (18.6%), *Klebsiella oxytoca*

(14.3%), and *Klebsiella pneumoniae* (11.9%). Our research findings can be compared to a study carried out in hospitals in India, *Staphylococcus aureus* also showed a relatively high occurrence (53%) (Mogasale *et al.*, 2021). In another study, *A. baumannii* and *K. pneumoniae* were the most common gram-negative isolates and had 89% and 81% of possible extensive drug resistance (P-XDR), respectively (Vaithiyam *et al.*, 2020). Among gram-positive pathogens *S. aureus*, 63% of isolates were MDR followed by P-XDR (22%) and only 15% showed non-multidrug-resistant (NMDR) pattern (Vaithiyam *et al.*, 2020). Furthermore, a study conducted in Tanzania also reported that *Staphylococcus aureus* (n=100; 22.8%), was the most commonly isolated bacteria. Seventy-eight (78) isolates of *Staphylococcus aureus* were obtained and from the antibiotic sensitivity test results, 27 (34.6%) were identified as methicillin-resistant *Staphylococcus aureus* (MRSA) (Moremi *et al.*, 2016). The resistance rate of *Klebsiella pneumoniae* to third-generation cephalosporins was 38.5% (Moremi *et al.*, 2016)

Increased levels of antibiotic resistance can be attributed to various causes: (i) there could be a direct relationship between the emergence of resistant strains and the magnitude of antibiotic consumption, (ii) the usage level of amoxicillin and ampicillin in Indonesia is getting to an alarming stage, (iii) the inappropriate doses use and therapy timing potentially increase resistance (Murniati, 2012). In a recent research on a serious infection of children in South-East Asia, 17% of blood culture isolates were not susceptible to ampicillin, penicillin, gentamicin, or a combination of these drugs (Frost *et al.*, 2018). In this study, overall antibiotic resistance was relatively low. As shown in Figure 2, with a cutting score of 50%, all the tested antibiotics still appear potent, except for penicillin. However, using these drugs needs to be based on the bacteria's sensitivity to antibiotics.

Resistance mechanisms can be transmitted from one bacterium to another either vertically, when inherited from relatives, or horizontally, through plasmids, which potentially transfer resistance among different species (Vázquez-López *et al.*, 2020). The primary cause of resistance is the misuse of antibiotics (Ghosh *et al.*, 2020). This consequently has an impact on treatment efforts, more difficulties and higher health costs. As reported by the World Health Organisation infections by resistant bacteria result in ineffective treatment, and increased risk of infection spread to others (WHO, 2015).

Conclusion

The most common type of infection observed in this study was pyoderma and the antibiotic treatment administered to most occurring cases of infection was amoxicillin. In addition, several groups of bacteria were resistant to multiple groups of antibiotics. According to the antibiotic sensitivity test result, the level of resistance from the highest percentage to the least is as follows: penicillin > ampicillin > tetracycline > cefuroxime > amikacin > gentamicin > cefotaxime > chloramphenicol > ceftriaxone > ceftazidime.

Acknowledgement

The authors are thankful to Pakem primary health care Yogyakarta and Microbiology Laboratory Faculty of Medicine, Public Health, and Nursing Universitas Gadjah Mada Yogyakarta for providing facilities to carry out this work. We also thank to Faculty of Pharmacy Universitas Gadjah Mada for financial support through Prime Research Grant.

References

Apsari, D. P., Dwicandra, N. M. O., & Jaelani, A. K. (2017). Pola Peresepan Antibiotik Pada Manajemen Faringitis Akut Dewasa Di Puskesmas. *Jurnal Endurance*, *2*(3):252. <https://doi.org/10.22216/jen.v2i3.2039>

Benjamin, W. (2019). Aktivitas Antibakteri Amoksisilin Terhadap Bakteri Gram Positif dan Bakteri Gram Negatif. *Jurnal Pijar Mipa*, *14*(3):189–191. <https://doi.org/10.29303/jpm.1029>

Cosgrove, S. E. & Carmeli, Y. (2003). The Impact of Antimicrobial Resistance on Health and Economic Outcomes. *36*:1433–1437

Estiningsih, D., Puspitasari, I. & Nuryastuti, T. (2016). Identifikasi Infeksi Multidrug-Resistant Organisms (MDRO) Pada Pasien Yang Dirawat di Bangsal Neonatal Intensive Care Unit (NICU) RSUP DR. Soeradji Tirtonegoro Klaten. *Jurnal Manajemen Dan Pelayanan Farmasi, Journal of Management and Pharmacy Practice*, *6*(3):243–248 <https://doi.org/10.22146/JMPF.351>

Fair, R. J. & Tor, Y. (2014). Perspectives in Medicinal Chemistry Antibiotics and Bacterial Resistance in the 21st Century. 25–64. <https://doi.org/10.4137/PMC.S14459>

Fenosa, A., Fusté, E., Ruiz, L., Veiga-Crespo, P., Vinuesa, T., Guallar, V., Villa, T. G. & Viñas, M. (2009). Role of TolC in *Klebsiella oxytoca* resistance to antibiotics. *Journal of Antimicrobial Chemotherapy*, *63*(4):668–674. <https://doi.org/10.1093/jac/dkp027>

Friedman, N. D., Temkin, E. & Carmeli, Y. (2016). The negative impact of antibiotic resistance. *Clinical*

Microbiology and Infection, *22*(5):416–422. <https://doi.org/10.1016/j.cmi.2015.12.002>

Frost, I., Laxminarayan, R., McKenna, N., Chai, S. & Joshi, J. (2018). Antimicrobial resistance and primary health care. World Health Organisation, 3–6. https://apps.who.int/iris/bitstream/handle/10665/326454/WHO-HIS-SDS-2018.56-eng.pdf?0Ahttps://www.who.int/docs/default-source/primary-health-care-conference/amr.pdf?sfvrsn=8817d5ba_2

Ghosh, D., Veeraraghavan, B., & Elangovan, R. (2020). cross Antibiotic Resistance and Epigenetics : More to It than Meets the Eye. *American Society for Microbiology*, *64*(2):1–14

Guo, Y., Song, G., Sun, M., Wang, J., & Wang, Y. (2020). Prevalence and Therapies of Antibiotic-Resistance in *Staphylococcus aureus*. In *Frontiers in Cellular and Infection Microbiology* *10*:107. Frontiers Media S.A. <https://doi.org/10.3389/fcimb.2020.00107>

Ho, J. & Ip, M. (2019). Antibiotic-Resistant Community-Acquired Bacterial Pneumonia. *Infectious Disease Clinics of North America*, *33*(4):1087–1103 <https://doi.org/10.1016/j.idc.2019.07.002>

Humaida, R. (2014). Strategy To Handle Resistance of Antibiotics. Rifka Humaida | Strategy To Handle Resistance of Antibiotics J MAJORITY, *3*:113

Igrejas, G., Zhang, J., da Silva Pranchevicius, M.-C., Ferreira, R. L., M da Silva, B. C., Rezende, G. S., Nakamura-Silva, R., Pitondo-Silva, A., Boni Campanini, E., A Brito, M. C., L da Silva, E. M., César de Melo Freire, C. & da Cunha, A. F. (2019). High Prevalence of Multidrug-Resistant *Klebsiella pneumoniae* harboring Several Virulence and β -Lactamase Encoding Genes in a Brazilian Intensive Care Unit. <https://doi.org/10.3389/fmicb.2018.03198>

Kemendes. (2011). Lampiran Peraturan Menteri Kesehatan Nomor 2406/Menkes/Per/Xii/2011 Tentang Pedoman Umum Penggunaan Antibiotik. Lampiran Peraturan Menteri Kesehatan Nomor 2406/Menkes/Per/Xii/2011 *Tentang Pedoman Umum Penggunaan Antibiotik*, *874*:35–38

Kemendes. (2017). Peraturan Menteri Kesehatan Republik Indonesia No. 27 Tentang Pedoman Pencegahan dan Pengendalian Infeksi di Fasilitas Pelayanan Kesehatan. 110265-110493

Kyriakidis, I., Vasileiou, E., Pana, Z. D. & Tragiannidis, A. (2021). Acinetobacter baumannii antibiotic resistance mechanisms. *Pathogens*, *10*(3):1–31. <https://doi.org/10.3390/pathogens10030373>

Laux, C., Peschel, A. & Krismer, B. (2019). *Staphylococcus aureus* Colonization of the human nose and interaction with other microbiome members. *Gram-Positive Pathogens*, *2*:723–730. <https://doi.org/10.1128/9781683670131.ch45>

Mahdani, W., Hayati, Z. & Yusriadi, T. (2020). Peta Distribusi dan Resistensi *Acinetobacter baumannii* dari Spesimen Klinik di RSUD Dr. Zainoel Abidin tahun 2018. *AVERROUS: Jurnal Kedokteran Dan Kesehatan Malikussaleh*, *6*(1):108. <https://doi.org/10.29103/averrous.v6i1.2666>

- Mogasale, V. V., Saldanha, P., Pai, V., Rekha, P. D. & Mogasale, V. (2021). A descriptive analysis of antimicrobial resistance patterns of WHO priority pathogens isolated in children from a tertiary care hospital in India. *Scientific Reports*, **11**(1):1–7. <https://doi.org/10.1038/s41598-021-84293-8>
- Moremi, N., Claus, H., & Mshana, S. E. (2016). Antimicrobial resistance pattern: A report of microbiological cultures at a tertiary hospital in Tanzania. *BMC Infectious Diseases*, **16**(1):1–7. <https://doi.org/10.1186/s12879-016-2082-1>
- Murniati. (2012). Tingkat Kepatuhan Pasien Tentang Penggunaan Antibiotika (Amoxicillin Dan Ampisilin) Di Puskesmas Tamalanrea Jaya Kota Makassar. *Jurnal Farmasi Sandi Karsa*, **6**(1):99–104
- Murray, C. J., Ikuta, K. S., Sharara, F., Swetschinski, L., Robles Aguilar, G., Gray, A., Han, C., Bisignano, C., Rao, P., Wool, E., Johnson, S. C., Browne, A. J., Chipeta, M. G., Fell, F., Hackett, S., Haines-Woodhouse, G., Kashef Hamadani, B. H., Kumaran, E.A.P., McManigal, B., ... Naghavi, M. (2022). Global burden of bacterial antimicrobial resistance in 2019: a systematic analysis. *The Lancet*, **399**(10325):629–655. [https://doi.org/10.1016/S0140-6736\(21\)02724-0](https://doi.org/10.1016/S0140-6736(21)02724-0)
- Siegel, J. D., Rhinehart, E., Jackson, M., Chiarello, L. & The Healthcare Infection Control Practices Advisory Committee. (2017). Management of Multidrug-resistant Organisms in Healthcare Setting. *Centers for Disease Control and Prevention*, 1–74
- Sinto, R. (2020). Peran Penting Pengendalian Resistensi Antibiotik pada Pandemi COVID-19. *Jurnal Penyakit Dalam Indonesia*, **7**(4):2020–2021
- Ur Rahman, S., Ali, T., Ali, I., Khan, N. A., Han, B. & Gao, J. (2018). The Growing Genetic and Functional Diversity of Extended Spectrum Beta-Lactamases. *BioMed Research International*, 2018. <https://doi.org/10.1155/2018/9519718>
- Vaithiyam, V. S., Rastogi, N., Ranjan, P., Mahishi, N., Kapil, A., Dwivedi, S. N., Soneja, M., Wig, N., & Biswas, A. (2020). Antimicrobial Resistance Patterns in Clinically Significant Isolates from Medical Wards of a Tertiary Care Hospital in North India. *Journal of Laboratory Physicians*, **12**(03):196–202. <https://doi.org/10.1055/s-0040-1721161>
- Vázquez-López, R., Solano-Gálvez, S. G., José, J., Vignon-Whaley, J., Andrés, J., Vaamonde, A., Andrés, L., Alonzo, P., Rivera Reséndiz, A., Muleiro Álvarez, M., Nabil Vega López, E., Franyuti-Kelly, G., Abelardo Álvarez-Hernández, D., Guzmán, V. M., Ernesto Juárez Bañuelos, J., Felix, J. M., Antonio González Barrios, J., & Fortes, T. B. (2020). *Acinetobacter baumannii* Resistance: A Real Challenge for Clinicians. *Antibiotics*, **205**. <https://doi.org/10.3390/antibiotics9040205>
- WHO. (2015). Global Action Plan on Antimicrobial Resistance. *Microbe Magazine*, **10**(9):354–355 <https://doi.org/10.1128/microbe.10.354.1>
- Zhen, X., Lundborg, C. S., Sun, X., Hu, X. & Dong, H. (2019). Economic burden of antibiotic resistance in ESKAPE organisms: A systematic review. In *Antimicrobial Resistance and Infection Control*, **8**(1):137 BioMed Central Ltd. <https://doi.org/10.1186/s13756-019-0590-7>